

Growth rate of small-scale dynamo at low magnetic Prandtl numbers

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2012 Phys. Scr. 86 018404

(<http://iopscience.iop.org/1402-4896/86/1/018404>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 193.166.207.137

The article was downloaded on 07/07/2012 at 18:23

Please note that [terms and conditions apply](#).

Growth rate of small-scale dynamo at low magnetic Prandtl numbers

Nathan Kleorin and Igor Rogachevskii

Department of Mechanical Engineering, Ben-Gurion University of the Negev, PO Box 653,
Beer-Sheva 84105, Israel
NORDITA, AlbaNova University Center, Roslagstullsbacken 23, SE-10691 Stockholm, Sweden
E-mail: gary@bgu.ac.il

Received 14 December 2011

Accepted for publication 24 May 2012

Published 4 July 2012

Online at stacks.iop.org/PhysScr/86/018404

Abstract

In this study, we discuss two key issues relating to a small-scale dynamo instability at low magnetic Prandtl numbers and large magnetic Reynolds numbers, namely: (i) the scaling for the growth rate of a small-scale dynamo instability in the vicinity of the dynamo threshold; (ii) the existence of the Golitsyn spectrum of magnetic fluctuations in small-scale dynamos. There are two different asymptotics for the small-scale dynamo's growth rate: in the vicinity of the threshold of the excitation of the small-scale dynamo instability, $\lambda \propto \ln(Rm/Rm^{cr})$, and when the magnetic Reynolds number is much larger than the threshold of the excitation of the small-scale dynamo instability, $\lambda \propto Rm^{1/2}$, where Rm^{cr} is the small-scale dynamo instability threshold in the magnetic Reynolds number Rm . We demonstrate that the existence of the Golitsyn spectrum of magnetic fluctuations requires a finite correlation time of the random velocity field. On the other hand, the influence of the Golitsyn spectrum on the small-scale dynamo instability is minor. This is the reason why it is so difficult to observe this spectrum in direct numerical simulations for the small-scale dynamo with low magnetic Prandtl numbers.

1. Introduction

Generation of a magnetic field by turbulent motions of conducting fluid is a fundamental mechanism of the magnetic fields observed in stars, galaxies and planets. There are different kinds of turbulent dynamos: large-scale and small-scale. The large-scale mean-field dynamo implies that the amplification of the magnetic field occurs at scales which are much larger than the maximum scale of the turbulent motion. This kind of dynamo includes: (i) the $\alpha\Omega$ and $\alpha^2\Omega$ dynamos caused by the combined action of the α effect and differential rotation (see, e.g. [1–5]); (ii) α^2 dynamo in helical turbulence; and (iii) the shear dynamos in non-helical turbulence [6–8].

On the other hand, generation of magnetic fluctuations occurs at scales which are smaller than the maximum scale of the turbulent motions (see, e.g. reviews [9–14]). Self-excitation of magnetic fluctuations with a zero mean magnetic field is called a small-scale dynamo. The mechanisms of the small-scale dynamo action are different depending on magnetic Prandtl numbers $Pm = \nu/\eta$, where ν is the kinematic viscosity of the fluid and η is the magnetic diffusion, due to the electrical conductivity

of the fluid. For large magnetic Prandtl numbers, the self-excitation of magnetic fluctuations is caused by the random stretching of the magnetic field by the smooth velocity fluctuations (see, e.g. [9, 10, 15–21]). This type of dynamo has been comprehensively studied in direct numerical simulations (DNS) of forced turbulence [22–25] and turbulent convection [26, 27]. The nature of the small-scale dynamo for low magnetic Prandtl numbers is different, e.g. it is driven by the inertial-range velocity fluctuations at the resistive scale. The small-scale dynamo at low magnetic Prandtl numbers has been studied analytically (see, e.g. [28–34]) for a Gaussian white-noise velocity field (so called the Kazantsev–Kraichnan model) and numerically (see, e.g. [14, 35, 36]) in a number of publications. Since the magnetic energy is not conserved, the second moment of the magnetic field has anomalous scalings [29, 37].

The small-scale dynamo instability is excited when the magnetic Reynolds number, Rm , is larger than the critical magnetic Reynolds number, Rm^{cr} . Analytical models based on the Kazantsev–Kraichnan model of a homogeneous, isotropic, non-helical and incompressible velocity field, yield $Rm^{cr} \approx 410$ at very low magnetic Prandtl numbers [29]. The compressibility of fluid flow causes a strong increase of the

critical magnetic Reynolds number at $Pm \ll 1$ (see [29]). A similar tendency has also recently been demonstrated in an analytical study [21] at large Prandtl numbers. DNSs of a small-scale dynamo in [14, 35, 36] of the Navier–Stokes turbulence show that Rm^{ct} is around 200 for small magnetic Prandtl numbers, and it is at three times larger than for the small-scale dynamo at large and moderate Prandtl numbers (see [22–24]). These DNS results at large and moderate Prandtl numbers are in agreement with different analytical models [10, 18, 21, 32].

The existence of the small-scale dynamo for a large number of turbulent spectra at large Prandtl numbers has been demonstrated [21]. When $Pm \sim 1$, the small-scale dynamo exists even in the regime of very large Mach numbers (see the DNS results in [25]). This study has also shown that, for low Mach numbers (~ 0.1), the ratio of the growth rate of turbulence driven by solenoidal and compressive forcing is about 30. However, for higher Mach numbers (~ 10), this ratio is about 2.

The small-scale dynamo action is different from the turbulent induction effect that causes the production of anisotropic magnetic fluctuations by a tangling of the mean magnetic field by these velocity fluctuations (see, e.g. [38–43]). This effect cannot be described in terms of the small-scale dynamo instability.

In spite of a number of studies of small-scale dynamos at low magnetic Prandtl numbers, there are some key questions that are the subject of discussions in the literature. One of them is related to scaling the growth rate λ of the small-scale dynamo instability at low magnetic Prandtl numbers in the vicinity of the dynamo threshold. Our analysis performed in this study, and even the numerical solution of the dynamo equations for a Gaussian white-noise velocity field obtained in [44], imply that there are two different asymptotics for the dynamo instability growth rate: (i) in the vicinity of the threshold of the excitation of the small-scale dynamo instability and (ii) far from the threshold of the small-scale dynamo instability.

Another issue studied here is related to the existence of the Golitsyn spectrum, $k^{-11/3}$, of magnetic fluctuations [38, 39] in the small-scale dynamo with low magnetic Prandtl numbers. This spectrum of magnetic fluctuations has been observed in the laboratory experiments [45, 46], in the large-eddy-simulations [47] and in the DNS [14, 36] of the small magnetic Prandtl numbers magnetohydrodynamic turbulence. In this study, we discuss the conditions for the existence of the Golitsyn spectrum.

The small-scale dynamo mechanism appears to be responsible for the random magnetic fields generated in the interstellar medium and in galaxy clusters [13, 14, 48–50]. A number of recent studies also pointed out the relevance of the small-scale dynamo in amplifying small seed fields in galaxies and the intergalactic medium (see, e.g. [51–55]). In particular, a DNS study [54] demonstrated that in the presence of turbulence, weak seed magnetic fields were amplified by the small-scale dynamo during the formation of the first stars. Strong magnetic fields were generated during the birth of the first stars in the universe, potentially modifying the mass distribution of these stars and influencing the subsequent cosmic evolution (see [54]). It was also noted [53] that the

small-scale dynamo was very efficient during the formation of the first stars and galaxies. During gravitational collapse, turbulence is created from accretion shocks, which may act to amplify weak magnetic fields in the protostellar cloud. Such turbulence is sub-sonic in the first star-forming minihalos, and highly supersonic in the first galaxies. It was concluded [53] that magnetic fields are significantly enhanced before the formation of a protostellar disk, where they may change the fragmentation properties of the gas and the accretion rate.

2. Governing equations

Let us study magnetic fluctuations with a zero mean magnetic field at low magnetic Prandtl numbers. In sections 2 and 3, we use the Kazantsev–Kraichnan model [28] of the δ -correlated-in-time random velocity field. Using this model allows us to get the analytical results for the growth rate of the small-scale dynamo instability. The results also remain valid for the velocity field with a finite correlation time if the second-order correlation functions of the magnetic field vary slowly, in comparison to the correlation time of the turbulent velocity field (see, e.g. [10, 56]). The two-point instantaneous correlation function of the magnetic field can be presented in the form

$$\langle b_i(t, \mathbf{x}) b_j(t, \mathbf{y}) \rangle = \tilde{W}(t, r) \delta_{ij} + \frac{r \tilde{W}'}{2} (\delta_{ij} - r_{ij}), \quad (1)$$

where $\tilde{W}(t, r) = \langle b_r(t, \mathbf{x}) b_r(t, \mathbf{y}) \rangle$ is the longitudinal correlation function, b_r is the component of magnetic field \mathbf{b} in the direction $\mathbf{r} = \mathbf{x} - \mathbf{y}$, $r_{ij} = r_i r_j / r^2$ and $\tilde{W}' = \partial \tilde{W} / \partial r$. This form of the second moment (1) corresponds to the condition $\nabla \cdot \mathbf{b} = 0$ and an assumption of the homogeneous and isotropic magnetic fluctuations. The equation for the function $\tilde{W}(r, t)$ derived in the framework of the Kazantsev–Kraichnan model of a homogeneous, isotropic, non-helical, incompressible and Gaussian white-noise velocity field, reads

$$\frac{\partial \tilde{W}(t, r)}{\partial t} = \frac{1}{m(r)} \tilde{W}'' + \mu(r) \tilde{W}' - \frac{\kappa(r)}{m(r)} \tilde{W}, \quad (2)$$

(see [28, 29]), where

$$\frac{1}{m(r)} = \frac{2}{Rm} + \frac{2}{3} [1 - F(r)], \quad \mu(r) = \frac{4}{mr} + \left(\frac{1}{m} \right)',$$

$$\kappa(r) = \frac{2m}{r} f'(r), \quad f(r) = F(r) + rF'/3,$$

and $Rm = u_0 \ell_0 / \eta \gg 1$ is the magnetic Reynolds number, u_0 is the characteristic turbulent velocity in the integral scale ℓ_0 and $F' = dF(r)/dr$. Hereafter equations are written in dimensionless variables: length and velocity are measured in units of ℓ_0 and u_0 . For a homogeneous, isotropic and non-helical (with zero mean helicity), incompressible turbulent fluid velocity field, the correlation function $\langle \tau u_i(\mathbf{x}) u_j(\mathbf{x} + \mathbf{r}) \rangle$ is given by

$$\langle \tau u_i(\mathbf{x}) u_j(\mathbf{y}) \rangle = \frac{1}{3} \left[F(r) \delta_{ij} + \frac{rF'}{2} (\delta_{ij} - r_{ij}) \right]. \quad (3)$$

The form of the continuous function $F(r)$ with different scalings in different ranges of scales is constructed using the following reasoning. The function $F(r) = 1 - \sqrt{Re} r^2$ is in the viscous range of scales: $0 \leq r \leq \ell_v/\ell_0$, while the function $F(r) = 1 - r^{4/3}$ is in the inertial range of scales: $\ell_v/\ell_0 < r < 1$. At the boundary of these ranges, $r = \ell_v/\ell_0$, these functions coincide, where $\ell_v = \ell_0/Re^{3/4}$ is the viscous scale and ℓ_0 is the integral scale of turbulence.

The solution of equation (2) can be obtained using an asymptotic analysis (see, e.g. [10, 29, 30]). This analysis is based on the separation of scales. In particular, the solutions of equation (2) with a variable mass have different regions with different functions $m(r)$, $\mu(r)$ and $\kappa(r)$. Solutions in these different regions and their derivatives can be matched at their boundaries. The results obtained by this asymptotic analysis are presented below.

3. Asymptotic behaviour of the growth rate of magnetic fluctuations

Let us now discuss the asymptotic behaviour of the growth rate of magnetic fluctuations with a zero mean for small magnetic Prandtl numbers. We sought a solution of equation (2) for the longitudinal correlation function of the magnetic field in the form: $\tilde{W}(t, r) = \exp(\lambda t) W(r)$. In the viscous range of scales, $0 \leq r \leq \ell_v/\ell_0$, the function $F(r) = 1 - \sqrt{Re} r^2$ and the equation for the function $W(r)$ is given by

$$r W'' + 4 W' + \frac{10}{3} Pr_m Re^{3/2} r W = 0, \quad (4)$$

where $W' = dW(r)/dr$. The solution of equation (4) is given by

$$W(r) = r^{-3/2} J_{3/2} \left(\sqrt{\frac{10 Pr_m}{3}} Re^{3/4} r \right) \approx 1 - \frac{Pr_m Re^{3/2}}{3} r^2, \quad (5)$$

(see [29]), where $J_\alpha(y)$ is the Bessel function of the first kind; we have taken into account that $W(r = 0) = 1$.

In the inertial range of scales, $\ell_v/\ell_0 < r < 1$, the function $F(r) = 1 - r^{4/3}$ and the equation for the function $W(r)$ is given by

$$\left(1 + \frac{1}{3} Rm r^{4/3}\right) W'' + \frac{4}{r} \left(1 + \frac{4}{9} Rm r^{4/3}\right) W' + Rm \left(\frac{52}{27} r^{-2/3} - \frac{\lambda}{2}\right) W = 0, \quad (6)$$

where λ is the growth rate of small-scale dynamo instability. In the range of scales, $\ell_v/\ell_0 < r \ll \ell_\eta/\ell_0$ the equation for the function $W(r)$ reads

$$r^2 W'' + 4r W' + \frac{52}{27} Rm r^{4/3} W = 0, \quad (7)$$

where $\ell_\eta = \ell_0/Rm^{3/4}$ is the resistive scale. The solution of equation (7) is given by

$$W(r) = r^{-3/2} J_{9/4} \left(\sqrt{\frac{13 Rm}{3}} r^{2/3} \right) \approx 1 - \frac{Rm}{3} r^{4/3}, \quad (8)$$

(see [29]). On the other hand, in the range of scales, $\ell_\eta/\ell_0 \ll r < 1$ the equation for the function $W(r)$ is given by

$$9r^2 W'' + 48r W' + \left(52 - \frac{27\lambda}{2} r^{2/3}\right) W = 0. \quad (9)$$

The solution of equation (9) is

$$W(r) = C r^{-13/6} K_\alpha \left(\sqrt{\frac{27\lambda}{2}} r^{1/3} \right), \quad (10)$$

(see [34]), where $K_\alpha(y)$ is the real part of the modified Bessel function (Macdonald function) with $\alpha = (i/2)\sqrt{39}$. This solution was chosen to be finite at large r , with a positively defined spectrum, and it has the following asymptotics at scales $\ell_\eta/\ell_0 \ll r \ll \lambda^{-3/2}$ (see [29]):

$$W(r) = A_1 r^{-13/6} \cos \left(\sqrt{\frac{13}{12}} \ln r + \varphi_0 \right), \quad (11)$$

and at scales $\lambda^{-3/2} \ll r \ll 1$ (see [32]):

$$W(r) = A_2 r^{-7/3} \exp \left(-\sqrt{\frac{27\lambda}{2}} r^{1/3} \right). \quad (12)$$

Here A_1 and A_2 are the constants which are proportional to the constant C .

In the range of scales $r \gg 1$, the turbulence is absent ($F \rightarrow 0$), $1/m = 2/3$, $\mu(r) = 4/mr$ and

$$W(r) = A_3 r^{-2} \left(\sqrt{\lambda} + r^{-1} \right) \exp(-\lambda r), \quad (13)$$

(see [29]), where A_3 is a constant.

The scaling for the growth rate of small-scale dynamo instability which is far from the threshold, is estimated as an inverse resistive time scale:

$$\lambda \sim \frac{u_\eta}{\ell_\eta} \sim \frac{u_0}{\ell_0} Rm^{1/2}, \quad (14)$$

(see [39]), where $u_\eta = (\varepsilon \ell_\eta)^{1/3}$ is the characteristic turbulent velocity at the resistive scale, $u_0 = (\varepsilon \ell_0)^{1/3}$ and ε is the dissipation rate of turbulent kinetic energy. For the scaling $\lambda \propto Rm^{1/2}$, the condition $\lambda^{1/2} r^{1/3} \gg 1$ implies $r \gg Rm^{-3/4}$. Note that this matching of the solutions (8), (12) and their derivatives at the boundary of their regions, yields the dynamo growth rate (14).

However, the scaling, $\lambda \propto Rm^{1/2}$, is not valid in the vicinity of the threshold of the dynamo instability. Indeed, in the vicinity of the threshold when $\lambda \rightarrow 0$, there is only one range of the solution of equation (9), i.e. $\lambda^{1/2} r^{1/3} \ll 1$. In this range of scales the solution of equation (9) is determined by equation (11). The matching of the solutions (8), (11) and (13) and their derivatives at the boundary their regions yields the following growth rate of the small-scale dynamo instability in the vicinity of the threshold:

$$\lambda = \beta \ln \left(\frac{Rm}{Rm^{cr}} \right), \quad (15)$$

(see appendix A), where $\beta = 4/3$ is the exponent of the scaling of the correlation function $F(r)$ (i.e. it is the exponent




Figure 1. The growth rate of small-scale dynamo instability versus $\ln(Rm/Rm^{cr})$ in the vicinity of the instability threshold: solid line corresponds to the scaling $\lambda = \beta \ln(Rm/Rm^{cr})$ and squares are the results of the numerical solution of the dynamo equation for $W(r)$ for the Kazantsev–Kraichnan model of velocity field with zero kinetic helicity in the inertial range of scales taken from figure 1 in [44].

of the turbulent diffusivity scaling). In figure 1, we plot the growth rate (15) of small-scale dynamo instability versus $\ln(Rm/Rm^{cr})$ in the vicinity of the threshold of small-scale dynamo instability. In the same figure we also show the numerical solution (squares) of the dynamo equation (2) performed in [44] (for the Kazantsev–Kraichnan model in an inertial range of scales of fluid motions for $Re \sim 10^8$, and for $Rm \leq 10^7$), which demonstrates a perfect agreement between the scaling (15), shown by the solid line, and the numerical solution of the dynamo equation.

Note that the solution of equation (9) determined by equation (11), is generally a fast oscillating function at $r \ll 1$. However, for the first mode with the maximum growth rate, the spectrum of the eigenfunction is positively defined (see [10]). Since this solution is only valid in the vicinity of the dynamo threshold, the second and higher modes are not excited. Therefore, the resulting spectrum of the eigenfunctions are always positively defined.

In this study, we only discuss the regime of small magnetic Prandtl numbers. In the case of large magnetic Prandtl numbers and large fluid Reynolds numbers, the dynamo growth rate far from the threshold is $\lambda \sim Re^{1/2}$, i.e. it is determined by the Kolmogorov time scale (see, e.g. [20, 21, 39]). On the other hand, for small magnetic Prandtl numbers the dynamo growth rate far from the threshold is determined by the resistive time scale.

4. Magnetic fluctuations with the Golitsyn spectrum

In this section, we study the effect of magnetic fluctuations with the Golitsyn spectrum, $k^{-11/3}$, [38, 39] on the small-scale dynamo with low magnetic Prandtl numbers. This spectrum can exist at the scales $\ell_v \leq r \leq \ell_\eta$. Our goal is to determine the longitudinal correlation function $W(r)$ that corresponds to the Golitsyn spectrum. To this end we use the induction equation for the instantaneous magnetic field $\mathbf{H}(t, \mathbf{x})$ in an

incompressible velocity field $\mathbf{v}(t, \mathbf{x})$:

$$\frac{\partial \mathbf{H}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{H} = (\mathbf{H} \cdot \nabla) \mathbf{v} + \eta \Delta \mathbf{H}. \quad (16)$$

We seek the solution of equation (16) in the following form:

$$\mathbf{H}(t, \mathbf{x}) = [\mathbf{B}(t) + \mathbf{b}(t, \mathbf{x})] \exp(\lambda t/2), \quad (17)$$

where $\mathbf{B}(t)$ is the magnetic field in scales which are much larger than the resistive scale ℓ_η , while $\mathbf{b}(t, \mathbf{x})$ is the magnetic field in the scales which are smaller than ℓ_η . We consider the magnetic dynamo regime, so that the total magnetic field \mathbf{H} grows in time exponentially with the growth rate $\lambda/2$. Next, we average equation (16) over the ensemble of fluctuations generated in the scales $\ell_\eta \ll \ell \ll \ell_0$, and subtract the obtained averaged equation from equation (16). This yields an equation for the magnetic field $\mathbf{b}(t, \mathbf{x})$:

$$\frac{\partial \mathbf{b}}{\partial t} = (\mathbf{B} \cdot \nabla) \mathbf{u} + (\eta \Delta - \lambda/2) \mathbf{b} + \mathbf{b}^N, \quad (18)$$

where $\mathbf{v}(t, \mathbf{x}) = \mathbf{V}(t) + \mathbf{u}(t, \mathbf{x})$, $\mathbf{V}(t)$ is the velocity field in the scales which are much larger than the resistive scale ℓ_η , while $\mathbf{u}(t, \mathbf{x})$ is the velocity field in the scales which are smaller than the resistive scale ℓ_η , and $\mathbf{b}^N = \nabla \times (\mathbf{u} \times \mathbf{b} - (\mathbf{u} \times \mathbf{b}))$ are the nonlinear terms. Equation (18) is written in the frame moving with the velocity $\mathbf{V}(t)$. Using equation (18) and the momentum equation for the velocity $\mathbf{u}(t, \mathbf{x})$ we derive equations for the second moments of the magnetic field $h_{ij}(\mathbf{k}) = \langle b_i(\mathbf{k}) b_j(-\mathbf{k}) \rangle$ and the cross helicity tensor $g_{ij}(\mathbf{k}) = \langle b_i(\mathbf{k}) u_j(-\mathbf{k}) \rangle$:

$$\begin{aligned} \frac{\partial h_{ij}(\mathbf{k})}{\partial t} = & -i(\mathbf{B} \cdot \mathbf{k}) [g_{ij}(\mathbf{k}) - g_{ji}(-\mathbf{k})] \\ & - (\eta k^2 + \lambda/2) h_{ij} + h_{ij}^N, \end{aligned} \quad (19)$$

$$\frac{\partial g_{ij}(\mathbf{k})}{\partial t} = i(\mathbf{B} \cdot \mathbf{k}) \tilde{f}_{ij}(\mathbf{k}) - (\eta k^2 + \lambda/2) g_{ij} + g_{ij}^N, \quad (20)$$

where $\tilde{f}_{ij}(\mathbf{k}) = \langle u_i(\mathbf{k}) u_j(-\mathbf{k}) \rangle$, $h_{ij}^N = \langle b_i^N(\mathbf{k}) b_j(-\mathbf{k}) \rangle + \langle b_i(\mathbf{k}) b_j^N(-\mathbf{k}) \rangle$ and $g_{ij}^N = -i(\mathbf{k} \cdot \mathbf{B}) h_{ij}(\mathbf{k}) + \langle b_i^N(\mathbf{k}) u_j(-\mathbf{k}) \rangle + \langle b_i(\mathbf{k}) u_j^N(-\mathbf{k}) \rangle$. Here u_i^N are the nonlinear terms in the momentum equation. Since we have already taken into account the exponential growth of the total magnetic field \mathbf{H} , we can drop the time derivatives in equations (19) and (20), because the characteristic times of the variations of the correlation functions h_{ij} and g_{ij} are much larger than the time λ^{-1} . Since we describe magnetic fluctuations in spatial scales which are smaller than the resistive scale ℓ_η , we may drop the nonlinear terms h_{ij}^N and g_{ij}^N in equations (19) and (20) in the case of large magnetic Reynolds numbers and low Prandtl numbers, because they are small in these scales. Therefore, equations (19) and (20) yield:

$$(\eta k^2 + \lambda/2)^2 \langle b_i(\mathbf{k}) b_j(-\mathbf{k}) \rangle = 2(\mathbf{B} \cdot \mathbf{k})^2 \langle u_i(\mathbf{k}) u_j(-\mathbf{k}) \rangle. \quad (21)$$

In the next step we introduce the normalized two-point correlation function $w(r)$ of the magnetic field which is

defined as follows:

$$w(r) = \frac{1}{\langle B^2 \rangle} \langle H_m(\mathbf{x}) H_m(\mathbf{y}) \rangle = 1 + \frac{1}{\langle B^2 \rangle} \langle b_m(\mathbf{x}) b_m(\mathbf{y}) \rangle, \quad (22)$$

where $w' = dw(r)/dr$ and $r = |\mathbf{x} - \mathbf{y}|$. We rewrite equation (21) in \mathbf{r} space using the inverse Fourier transformation (i.e. we use the following transformation $ik_i \rightarrow \nabla_i$). This yields the following equation for the normalized two-point correlation function of the magnetic field:

$$\left(\frac{d^2}{dr^2} + \frac{2}{r} \frac{d}{dr} - a^2 \right)^2 w(r) = -\frac{2}{3} Rm^2 \left(\frac{d^2 \tilde{f}(r)}{dr^2} + \frac{2}{r} \frac{d\tilde{f}(r)}{dr} \right) + 3a^4, \quad (23)$$

where $a = (\lambda Rm/2)^{1/2}$, $\tilde{f}(r) = \tilde{f}_{mm}(r)$ and $\tilde{f}_{ij}(r)$ is the two-point correlation function of the velocity fluctuations written in \mathbf{r} space. Equation (23) is written in dimensionless variables: length and velocity are measured in units of ℓ_0 and u_0 , the growth rate of the magnetic fluctuations λ is measured in units of u_0/ℓ_0 and the magnetic field is measured in units of B_0 . We also take into account that $\langle (\mathbf{B} \cdot \nabla)^2 \tilde{f} \rangle = (1/3) (\tilde{f}'' + 2\tilde{f}'/r)$.

The solution of equation (23), which satisfies the following boundary conditions: $w(r=0) = 3$ and $w'(r=0) = w''(r=0) = 0$, reads

$$w(r) = 3 - \frac{3C_1}{4a^3 r} [(ar - 1) \exp(ar) + (ar + 1) \exp(-ar)] + \frac{55 Rm^2}{2(3^6) a^{11/3} r} \left[(3ar - 5) \exp(ar) \gamma(2/3, ar) + (3ar + 5) \exp(-ar) \gamma(2/3, -ar) \right], \quad (24)$$

where $w' = dw(r)/dr$, C_1 is a free constant and $\gamma(\beta, x) = \beta^{-1} x^\beta \exp(-x) M(1, 1 + \beta, x)$ is the incomplete gamma function which is related to the confluent hypergeometric function $M(a, b, x)$. When $ar \ll 1$, equation (24) for the two-point correlation function $w(r)$ is given by

$$w(r) = 3 - C_1 r^2 + \frac{1}{12} Rm^2 r^{8/3} \left[1 + \frac{9}{238} Rm \lambda r^2 \right]. \quad (25)$$

The function $w(r)$ is related to the longitudinal correlation function $W(r)$, i.e. $w(r) = 3W(r) + rW'(r)$. Equation (26), rewritten for the longitudinal correlation function $W(r)$, reads

$$W(r) = 1 - \frac{Rm Re^{1/2}}{3} r^2 + \frac{1}{68} Rm^2 r^{8/3} \left[1 + \frac{9}{322} Rm \lambda r^2 \right], \quad (26)$$

where $\ell_v/\ell_0 \leq r \leq \ell_\eta/\ell_0$, the constant $C_1 \approx (5/3) Rm Re^{1/2}$ is determined by the matching of functions $W(r)$ determined by equations (5) and (26) at the point $r = \ell_v/\ell_0$. The scaling $W(r) \propto Rm^2 r^{8/3}$ corresponds to the Golitsyn spectrum $M(k) \propto B_0^2 \eta^{-2} \varepsilon^{2/3} k^{-11/3}$ [38], where ε is the rate of

dissipation of the turbulent kinetic energy and $M(k) = (2/\pi) \int_0^\infty kr \sin(kr) w(r) dr$. It follows from the latter equation that the exponent q in the spectrum function $M(k) \propto k^{-q}$ and the exponent p in the scaling $W(r) \propto r^p$ are related as follows: $q = p + 1$.

For small Prandtl numbers the constant $\tilde{C}_1 = Rm Re^{1/2}/3$ is larger than $\tilde{C}_2 = Rm^2/68$, and since $r \ll 1$, the first and second terms in the right hand side of equation (26) dominate the behaviour of $W(r)$. This estimate implies that the third term in the right hand side of equation (26) resembling the Golitsyn spectrum, is negligible. This is the reason why the influence of the Golitsyn spectrum on the small-scale dynamo instability is minor. That is why it is so difficult to observe the Golitsyn spectrum in DNS for the small-scale dynamo with low magnetic Prandtl numbers.

Note that the existence of the Golitsyn spectrum of magnetic fluctuations requires a finite correlation time of the random velocity field, i.e. the solution for the small-scale dynamo with the Golitsyn spectrum does not exist in the framework of the Kraichnan–Kazantsev model of the delta-correlated-in-time turbulent velocity field (see appendix B). Indeed, for the derivation of equation (23) we did not use the assumption about the delta-correlated-in-time turbulent velocity field. One of the indications of the finite correlation time of the random velocity field is already seen in equation (23), where the high-order spatial derivatives arise. The Kazantsev–Kraichnan model yields the dynamo equation with spatial derivatives not higher than the second-order spatial derivatives. On the other hand, it is well-known that even a small yet finite correlation time of the random velocity field causes the appearance of the higher-order spatial derivatives in the dynamo equations (see, e.g. [10, 56]).

The requirement of the finite correlation time of the random velocity field for the correct description of the tangling magnetic fluctuations which have the Golitsyn spectrum, also follows from the dimensional arguments. Indeed, the main balance in the induction equation for the magnetic fluctuations $(\mathbf{B} \cdot \nabla) \mathbf{u} \sim \eta \Delta \mathbf{b}$ which yields the Golitsyn spectrum, can be rewritten in the following form: $\langle \mathbf{b}^2 \rangle \sim \tau^2(\ell) [(\mathbf{u}^2)]^2 \mathbf{B}^2/\eta^2$. The latter equation implies the requirement of the finite correlation time of the random velocity field for the correct description of the tangling magnetic fluctuations. Similar arguments are also valid for the k^{-1} spectrum of the magnetic fluctuations generated by the tangling mechanism at low magnetic Prandtl numbers in the scales $\ell_\eta \ll \ell \ll \ell_0$ (see [40–43]).

We stress again that both magnetic fields, $\mathbf{B}(t)$ and $\mathbf{b}(t, \mathbf{x})$, are small-scale fields (in scales which are much less than the integral scale ℓ_0 of turbulence). In particular, $\mathbf{B}(t)$ is the magnetic field in the scales $\ell_\eta \ll \ell \ll \ell_0$, while $\mathbf{b}(t, \mathbf{x})$ is the magnetic field in the scales $\ell_v \ll \ell \ll \ell_\eta$. These fields belong to the same mode generated by the same small-scale dynamo mechanism. In this section, we used two magnetic fields, $\mathbf{B}(t)$ and $\mathbf{b}(t, \mathbf{x})$, to describe the interaction of the magnetic fields of different scales by the tangling of the field $\mathbf{B}(t)$ of the velocity fluctuations which produces additional anisotropic magnetic fluctuations with the Golitsyn spectrum. The latter mechanism is the turbulent magnetic induction that is different from the small-scale dynamo.

5. Conclusions

In this study, we investigated some key issues of small-scale dynamos in a random velocity field with large fluid Reynolds numbers, a zero mean magnetic field and low magnetic Prandtl numbers. Contrary to the claim in [44], there are two different asymptotics for the dynamo growth rate: in the vicinity of the threshold of the excitation of the dynamo instability ($\lambda \propto \ln(Rm/Rm^{cr})$) and far from the dynamo threshold ($\lambda \propto Rm^{1/2}$). The influence of the Golitsyn spectrum on the small-scale dynamo instability is minor, and this spectrum of magnetic fluctuations requires a finite correlation time of the random velocity field. However, the Golitsyn spectrum of magnetic fluctuations does exist in a small-scale turbulence at low magnetic Prandtl numbers with an imposed constant large-scale magnetic field.

Acknowledgments

We are grateful to Alexander Schekochihin for the stimulating discussions which initiated our study of small-scale dynamo with the Golitsyn spectrum. We acknowledge the NORDITA dynamo program of 2011 for providing a stimulating scientific atmosphere. This work was supported in part by COST Action MP0806, by the European Research Council under the Atmospheric Research Project no. 227915 and by a grant from the Government of the Russian Federation under contract no. 11.G34.31.0048.

Appendix A. Growth rate of small-scale dynamo instability in the vicinity of the dynamo threshold

Let us obtain the scaling for the growth rate of the small-scale dynamo instability in the vicinity of the dynamo threshold. In the range of scales, $\ell_v/\ell_0 < r \ll \ell_\eta/\ell_0$, the function rW'/W is given by

$$\frac{rW'}{W} = -\frac{4}{9} Rm r^{4/3}, \quad (\text{A.1})$$

(see equation (8)), while in the range of scales, $\ell_\eta/\ell_0 \ll r < 1$ the function rW'/W is

$$\frac{rW'}{W} = -\frac{13}{6} - \sqrt{\frac{13}{12}} \tan\left(\sqrt{\frac{13}{12}} \ln r + \varphi_0\right), \quad (\text{A.2})$$

(see equation (11)). On the other hand, in the range of scales $r \gg 1$ the function rW'/W is

$$\frac{rW'}{W} = -2 - \sqrt{\lambda} r - \frac{1}{1 + \sqrt{\lambda} r} \approx -(3 + \lambda r^2), \quad (\text{A.3})$$

(see equation (13)), where we have taken into account that in the vicinity of the dynamo threshold $\lambda \rightarrow 0$ and $\lambda^{1/2} r \ll 1$.

Matching of the functions rW'/W determined by equations (A.1) and (A.2) at the point $r = \ell_\eta/\ell_0$ yields the following equation:

$$\tan\left(\frac{\sqrt{39}}{8} \ln Rm - \varphi_0\right) = \frac{31}{3\sqrt{39}}. \quad (\text{A.4})$$

This equation determines the function $\varphi_0(Rm)$. Now we define the function $\varphi_0^{cr} = \varphi_0(Rm = Rm^{cr})$, where Rm^{cr} is the threshold for the excitation of the magnetic fluctuations. It follows from this equation that

$$\varphi_0 - \varphi_0^{cr} = \frac{\sqrt{39}}{8} \ln\left(\frac{Rm}{Rm^{cr}}\right). \quad (\text{A.5})$$

Matching of the functions rW'/W determined by equations (A.2) and (A.3) at the point $r = 1$ yields

$$\tan \varphi_0 = \sqrt{\frac{12}{13}} \left(\frac{5}{6} + \lambda\right). \quad (\text{A.6})$$

It follows from this equation that

$$\lambda = \frac{32}{3\sqrt{39}} (\varphi_0 - \varphi_0^{cr}), \quad (\text{A.7})$$

where we have also taken into account that in the vicinity of the dynamo threshold $\lambda \rightarrow 0$. Combining equations (A.5) and (A.7), we obtain the following scaling for the growth rate of small-scale dynamo instability in the vicinity of the dynamo threshold: $\lambda = (4/3) \ln(Rm/Rm^{cr})$.

Appendix B. Tangling magnetic fluctuations in the delta-correlated-in-time velocity field

The technique of path integrals for the delta-correlated-in-time velocity field allows us to derive the equation for the second-order correlation function, $h_{ij} = \langle b_i(t, \mathbf{x}) b_j(t, \mathbf{y}) \rangle$:

$$\frac{\partial h_{ij}}{\partial t} = [\hat{L}_{ik}(\mathbf{x}) \delta_{js} + \hat{L}_{js}(\mathbf{y}) \delta_{ik} + \hat{M}_{ijks}] h_{ks} + I_{ij}, \quad (\text{B.1})$$

(see for details [29]), where the turbulent component of magnetic field is $\mathbf{b}(t, \mathbf{x}) = \mathbf{H}(t, \mathbf{x}) - \mathbf{B}(t)$,

$$\hat{L}_{ij} = \frac{1}{3} \left(1 + \frac{3}{Rm}\right) \delta_{ij} \frac{\partial}{\partial x_p} \frac{\partial}{\partial x_p}, \quad (\text{B.2})$$

$$\begin{aligned} \frac{1}{2} \hat{M}_{ijks} &= \delta_{ik} \delta_{js} f_{mn} \frac{\partial^2}{\partial x_m \partial y_n} - \delta_{ik} \frac{\partial f_{mj}}{\partial y_s} \frac{\partial}{\partial x_m} - \delta_{js} \frac{\partial f_{in}}{\partial x_k} \frac{\partial}{\partial y_n} \\ &+ \frac{\partial^2 f_{ij}}{\partial x_k \partial y_s}, \end{aligned} \quad (\text{B.3})$$

$$I_{ij} = B_k B_s \frac{\partial^2 f_{ij}}{\partial x_k \partial y_s}, \quad (\text{B.4})$$

and $f_{mn} = \langle \tau u_m(\mathbf{x}) u_n(\mathbf{y}) \rangle$. Multiplying equation (B.1) by $r_i r_j / r^2$ we arrive at the equation for the correlation function $\tilde{W}(r, t)$:

$$\frac{\partial \tilde{W}(r, t)}{\partial t} = \frac{1}{m(r)} \tilde{W}'' + \mu(r) \tilde{W}' - \frac{\kappa(r)}{m(r)} \tilde{W} + I, \quad (\text{B.5})$$

where $I = B^2 (F'' + 4F'/r)/3$. In the inertial range the source term is $I = 52B^2 r^{-2/3}/27$. This source term yields the following scaling of the correlation function $W(r) \propto r^{4/3}$ in

the range of scales, $\ell_v/\ell_0 < r \ll \ell_\eta/\ell_0$. This implies that the scaling of magnetic fluctuations caused by the tangling of the large-scale magnetic field by the delta-correlated-in-time velocity field coincides with the scaling of turbulent magnetic diffusion $F(r) \propto r^{4/3}$. In Fourier space this corresponds to the $k^{-7/3}$ spectrum of magnetic fluctuations. This implies that the Golitsyn spectrum, $k^{-11/3}$, of magnetic fluctuations cannot be described in terms of the delta-correlated-in-time velocity field.

References

- [1] Moffatt H K 1978 *Magnetic Field Generation in Electrically Conducting Fluids* (New York: Cambridge University Press)
- [2] Parker E 1979 *Cosmical Magnetic Fields* (New York: Oxford University Press)
- [3] Krause F and Rädler K H 1980 *Mean-Field Magnetohydrodynamics and Dynamo Theory* (Oxford: Pergamon)
- [4] Zeldovich Ya B, Ruzmaikin A A and Sokoloff D D 1983 *Magnetic Fields in Astrophysics* (New York: Gordon and Breach)
- [5] Ruzmaikin A, Shukurov A M and Sokoloff D D 1988 *Magnetic Fields of Galaxies* (Dordrecht: Kluwer)
- [6] Rogachevskii I and Kleorin N 2003 *Phys. Rev. E* **68** 036301
- [7] Brandenburg A 2005 *Astrophys. J.* **625** 539
- [8] Yousef T A, Heinemann T, Schekochihin A A, Kleorin N, Rogachevskii I, Iskakov A B, Cowley S C and McWilliams J C 2008 *Phys. Rev. Lett.* **100** 184501
- [9] Zeldovich Ya B, Molchanov S A, Ruzmaikin A A and Sokoloff D D 1988 *Sov. Sci. Rev. C* **7** 1
- [10] Zeldovich Ya B, Ruzmaikin A A and Sokoloff D D 1990 *The Almighty Chance* (Singapore: World Scientific)
- [11] Childress S and Gilbert A 1995 *Stretch, Twist, Fold: The Fast Dynamo* (Berlin: Springer)
- [12] Kulsrud R M and Anderson S W 1992 *Astrophys. J.* **396** 606
- [13] Brandenburg A and Subramanian K 2005 *Phys. Rep.* **417** 1
- [14] Schekochihin A A, Iskakov A B, Cowley S C, McWilliams J C, Proctor M R E and Yousef T A 2007 *New J. Phys.* **9** 300
- [15] Batchelor G K 1950 *Proc. R. Soc. A* **201** 405
- [16] Moffatt H K and Saffman P G 1963 *Phys. Fluids* **7** 155
- [17] Zeldovich Y B, Ruzmaikin A A, Molchanov S A and Sokoloff D D 1984 *J. Fluid Mech.* **144** 1
- [18] Kleorin N and Rogachevskii I 1994 *Phys. Rev. E* **50** 493
- [19] Kleorin N, Rogachevskii I and Eviatar A 1994 *J. Geophys. Res. A* **99** 6475
- [20] Subramanian K 1998 *Mon. Not. R. Astron. Soc.* **294** 718
- [21] Schober J, Schleicher D, Federrath Ch, Klessen R and Banerjee R 2012 *Phys. Rev. E* **85** 026303
- [22] Schekochihin A A, Cowley S C, Taylor S F, Maron J L and McWilliams J C 2004 *Astrophys. J.* **612** 276
- [23] Haugen N E L, Brandenburg A and Dobler W 2004 *Phys. Rev. E* **70** 016308
- [24] Schekochihin A A, Haugen N E L, Brandenburg A, Cowley S C, Maron J L and McWilliams J C 2005 *Astrophys. J.* **625** L115
- [25] Federrath C, Chabrier G, Schober J, Banerjee R, Klessen R S and Schleicher D R G 2011 *Phys. Rev. Lett.* **107** 114504
- [26] Brandenburg A, Jennings R L, Nordlund A, Rieutord M, Stein R F and Tuominen I 1996 *J. Fluid Mech.* **306** 325
- [27] Cattaneo F 1999 *Astrophys. J.* **515** L39
- [28] Kazantsev A P 1968 *Sov. Phys.—JETP* **26** 1031
- [29] Rogachevskii I and Kleorin N 1997 *Phys. Rev. E* **56** 417
- [30] Rogachevskii I and Kleorin N 1999 *Phys. Rev. E* **59** 3008
- [31] Vincenzi D 2002 *J. Stat. Phys.* **106** 1073
- [32] Boldyrev S and Cattaneo F 2004 *Phys. Rev. Lett.* **92** 144501
- [33] Celani A, Mazzino A and Vincenzi D 2006 *Proc. R. Soc. A* **462** 137
- [34] Arponen H and Horvai P 2007 *J. Stat. Phys.* **129** 205
- [35] Iskakov A B, Schekochihin A A, Cowley S C, McWilliams J C and Proctor M R E 2007 *Phys. Rev. Lett.* **98** 208510
- [36] Brandenburg A 2011 *Astrophys. J.* **741** 92
- [37] Vergassola M 1996 *Phys. Rev. E* **53** 3021
- [38] Golitsyn G S 1960 *Sov. Phys.—Dokl.* **5** 536
- [39] Moffatt K 1961 *J. Fluid Mech.* **11** 625
- [40] Ruzmaikin A A and Shukurov A M 1982 *Astrophys. Space Sci.* **82** 397
- [41] Kleorin N, Rogachevskii I and Ruzmaikin A 1990 *Sov. Phys.—JETP* **70** 878
- [42] Kleorin N and Rogachevskii I 1994 *Phys. Rev. E* **50** 2716
- [43] Kleorin N, Mond M and Rogachevskii I 1996 *Astron. Astrophys.* **307** 293
- [44] Malyshekin L M and Boldyrev S 2010 *Phys. Rev. Lett.* **105** 215002
- [45] Odier P, Pinton J-F and Fauve S 1998 *Phys. Rev. E* **58** 7397
- [46] Bourgoin M *et al* 2002 *Phys. Fluids* **14** 3046
- [47] Ponty Y, Politano H and Pinton J-F 2004 *Phys. Rev. Lett.* **92** 144503
- [48] Beck R, Brandenburg A, Moss D, Shukurov A and Sokoloff D 1996 *Annu. Rev. Astron. Astrophys.* **34** 155
- [49] Subramanian K, Shukurov A and Haugen N E L 2006 *Mon. Not. R. Astron. Soc.* **366** 1437
- [50] Enßlin T A and Vogt C 2006 *Astron. Astrophys.* **453** 447
- [51] Ryu D, Kang H, Cho J and Das S 2008 *Science* **320** 909
- [52] Arshakian T G, Beck R, Krause M and Sokoloff D 2009 *Astron. Astrophys.* **494** 21
- [53] Schleicher D R G, Banerjee R, Sur S, Arshakian T G, Klessen R S, Beck R and Spaans M 2010 *Astron. Astrophys.* **522** A115
- [54] Sur S, Schleicher D R G, Banerjee R, Federrath Ch and Klessen R S 2010 *Astrophys. J. Lett.* **721** L134
- [55] Turk M J, Oishi J S, Abel T and Bryan G L 2012 *Astrophys. J.* **745** 154
- [56] Kleorin N, Rogachevskii I and Sokoloff D 2002 *Phys. Rev. E* **65** 036303